

COLORADO ASTHMA CARE PLAN AND MEDICATION ORDER FOR SCHOOL AND CHILD CARE SETTINGS

PARENT/GUARDIAN COMPLETE AND SIGN:

School/grade: _____
 Child Name: _____ Birthdate: _____
 Parent/Guardian Name: _____ Phone: _____
 Healthcare Provider Name: _____ Phone: _____
 Triggers: ☐ Weather (cold air, wind) ☐ Illness ☐ Exercise ☐ Smoke ☐ Dust ☐ Pollen ☐ Other: _____
☐ Life threatening allergy, specify: _____

I give permission for school personnel to share this information, follow this plan, administer medication and care for my child/youth, and if necessary, contact our healthcare provider. I assume full responsibility for providing the school/program prescribed medication and supplies, and to comply with board policies, if applicable. I am aware 911 may be called if a quick relief inhaler is not at school and my child/youth is experiencing symptoms. I approve this care plan for my child/youth.

PARENT SIGNATURE		DATE	NURSE/CCHC SIGNATURE		DATE
HEALTHCARE PROVIDER COMPLETE ALL ITEMS, SIGN AND DATE:			QUICK RELIEF (RESCUE) MEDICATION: ☐ Albuterol ☐ Other: _____ Common side effects: ☐ heart rate, tremor ☐ Have child use spacer with inhaler. Controller medication used at home: _____		
IF YOU SEE THIS:			DO THIS:		
GREEN ZONE: No Symptoms Pretreat	- No current symptoms - Doing usual activities		Pretreat strenuous activity: ☐ Not required ☐ Routine ☐ Student/Parent request Give QUICK RELIEF MED 10-15 minutes before activity: ☐ 2 puffs ☐ 4 puffs ☐ Repeat in 4 hours, if needed for additional physical activity. <i>If child is currently experiencing symptoms, follow YELLOW ZONE.</i>		
	YELLOW ZONE: Mild symptoms - Trouble breathing - Wheezing - Frequent cough - Complains of tight chest - Not able to do activities, but talking in complete sentences - Peak flow: _____ & _____		1. Stop physical activity. 2. Give QUICK RELIEF MED: ☐ 2 puffs ☐ 4 puffs 3. Stay with child/youth and maintain sitting position. 4. REPEAT QUICK RELIEF MED, if not improving in 15 minutes: ☐ 2 puffs ☐ 4 puffs 5. Child/youth may go back to normal activities, once symptoms are relieved. 6. Notify parents/guardians and school nurse. <i>If symptoms do not improve or worsen, follow RED ZONE.</i>		
	RED ZONE: EMERGENCY Severe Symptoms - Coughs constantly - Struggles to breathe - Trouble talking (only speaks 3-5 words) - Skin of chest and/or neck pull in with breathing - Lips/fingernails gray or blue ↓ Level of consciousness - Peak flow < _____		1. Give QUICK RELIEF MED: ☐ 2 puffs ☐ 4 puffs ▪ Refer to anaphylaxis plan, if child/youth has life-threatening allergy. 2. Call 911 and inform EMS the reason for the call. 3. Stay with child/youth. Remain calm, encouraging slower, deeper breaths. 4. Notify parents/guardians and school nurse. 5. If symptoms do not improve, REPEAT QUICK RELIEF MED: ☐ 2 puffs ☐ 4 puffs every 5 minutes until EMS arrives. <i>School personnel should not drive student to hospital.</i>		
PROVIDER INSTRUCTIONS FOR QUICK RELIEF INHALER USE: CHECK APPROPRIATE BOX(ES) ☐ Student needs supervision or assistance to use inhaler. Student will not self-carry inhaler. ☐ Student understands proper use of asthma medications, and in my opinion, <u>can carry and use his/her inhaler at school independently with approval from school nurse and completion of contract.</u> ☐ Student will notify school staff after using quick relief inhaler, if symptoms do not improve with use.					
HEALTH CARE PROVIDER SIGNATURE		PRINT PROVIDER NAME	DATE	FAX	PHONE

Copies of plan provided to: ☐ Teacher(s) ☐ PhysEd/Coach ☐ Principal ☐ Main Office ☐ Bus Driver Other _____



COLORADO
Department of Education

Revised: March 2018